



A comment to the talk by E. Seiler

Salvador Miracle-Sole, Giovanni Gallavotti, Francesco Guerra

► To cite this version:

Salvador Miracle-Sole, Giovanni Gallavotti, Francesco Guerra. A comment to the talk by E. Seiler. Conference of the International Association of Mathematical Physics, 1977, Rome, Italy. pp.436-438. hal-00707916

HAL Id: hal-00707916

<https://hal.science/hal-00707916>

Submitted on 13 Jun 2012

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

A COMMENT TO THE TALK BY E. SEILER

G. Gallavotti, F. Guerra, S. Miracle-Solé

KEY WORDS: Lattice gauge theories; Wegner model; Wilson parameter.

The formula given by Wegner may be rigorously proven as follows (for instance).

Let Σ_L be a square surface with side L laying on a coordinate plane of the lattice \mathbf{Z}^3 on which an Ising model is resting.

Put for every pair of nearest neighbours $f = (i, j)$:

$$\sigma_f = \sigma_i \sigma_j$$

and let

$$\begin{aligned} \beta_f &= \beta^* & \text{if } f \cap \Sigma_L &= \emptyset \\ \beta_f &= -\beta^* & \text{if } f \cap \Sigma_L &\neq \emptyset \end{aligned}$$

We wish to show that:

$$\lim_{\Lambda \uparrow \mathbf{R}^3} \frac{Z_\Lambda}{Z_\Lambda^0} = \frac{\sum_\sigma \exp \sum_{f \subset \Lambda} \beta_f^* \sigma_f}{\sum_\sigma \exp \sum_{f \subset \Lambda} \beta^* \sigma_f} = \exp(O(\beta^4))L$$

This is an immediate consequence of the Gruber-Kunz cluster expansion which in our case may be performed as follows (if \mathcal{B}_Λ = set of subset of “bonds”, i.e., sets of n.n. pairs) :

$$\begin{aligned} Z_\Lambda &= \sum_\sigma \prod_f \left(1 + (e^{\beta^* \sigma_f} - 1) \right) = \sum_{F \subset \mathcal{B}_\Lambda} \sum_\sigma \prod_{f \in F} (e^{\beta^* \sigma_f} - 1) \\ &= 2^{|\Lambda|} \sum_{\substack{x_1, \dots, x_k \\ x_i \cap x_j = \emptyset, i \neq j}} \prod_{i=1}^k \zeta(x_i) \end{aligned}$$

where x_1, \dots, x_k are the “connected components” of the family F of bonds and

$$\zeta(x) = \sum_{\sigma_x} 2^{-|x|} \prod_{f \in x} (e^{\beta^* \sigma_f} - 1)$$

The logarithm of the sum expressing Z_Λ can be expanded into a series of products of the ζ ’s and it is well known that only “connected diagrams”

appear in this expansion. More precisely let $N_\Lambda =$ set of the families $\Gamma = (x_1, \dots, x_p)$ of connected components such that if $x_i, x_j \in \Gamma$ then there is a chain $i_1 = i, i_2, \dots, i_p = j$ for which $x_{i_k} \cap x_{i_{k+1}} \neq \emptyset$, $k = 1, \dots, p-1$.

The elements of Γ may contain several times the same x : so we may represent Γ as $(x_1^{n_1}, \dots, x_p^{n_p})$ where $n_1, \dots, n_p > 0$ are the (integer) multiplicities and $x_i \neq x_j$. Let $|\Gamma| = n_1 + \dots + n_p$. Gruber and Kunz results read in this case

$$Z_\Lambda = 2^{|\Lambda|} \exp \sum_{\Gamma \in N_\Lambda} \phi^T(\Gamma) \zeta(\Gamma) \quad (1)$$

provided the series converges absolutely, where if $\Gamma = (x_1^{n_1}, \dots, x_p^{n_p})$:

$$\zeta(\Gamma) = \prod_{i=1}^p \zeta(x_i)^{n_i} \quad (2)$$

and the numbers $\phi^T(\Gamma)$ are certain combinatorial constants (independent of Λ if $\Gamma \in N_\Lambda$) verifying

$$|\phi^T(\Gamma)| \leq |\Gamma|^{-1} \quad (3)$$

An expansion similar to the (1) above can be made for the Z_Λ^0 and calling $\zeta^0(x)$ the corresponding ζ 's

$$Z_\Lambda / Z_\Lambda^0 = \langle \exp -2\beta^* \sum_{f \in \Sigma_\Lambda} \sigma_f \rangle_0 = \exp \sum_{\Gamma \cap \Sigma_L \neq \emptyset} \phi^T(\Gamma) (\zeta(\Gamma) - \zeta^0(\Gamma))$$

since $\zeta(\Gamma) = \zeta^0(\Gamma)$ if $\Gamma \cap \Sigma_L = \emptyset$ (provided the series in (1) for Z_Λ and the analogous for Z_Λ^0 converge).

It is obvious, however, that


$$|\zeta(\Gamma)| \leq |e^{\beta^*} - 1| = \xi(\Gamma), \quad |\zeta^0(\Gamma)| \leq \xi(\Gamma)$$

which implies the convergence of the series (because of (2)) if β^* is small and also (since Γ is connected):

$$\sum_{\Gamma \ni 0} |\phi^T(\Gamma)| \xi(\Gamma) \leq O(\beta^*)^\ell$$

if β^* is small.

Finally remark that the Γ 's for which $\zeta(\Gamma) \neq \zeta^0(\Gamma)$ are those which contain at least one x containing a closed circuit intersecting Σ_L an odd number of times (recall that x is a set of bonds and therefore it may contain "closed circuits").



The convergence of the series implies that the lowest order dominates which corresponds to the sum

$$\exp \sum_{\gamma \in \partial \Sigma_L} \zeta(\gamma) \phi^T(\gamma)$$

where γ is the set of four pairs of n.n. forming a square as 1234 in the picture.

The $\phi^T(\gamma)$ for such a configuration is +1 and

$$\begin{aligned} \zeta(f) &= \sum_{\sigma_1, \sigma_2, \sigma_3, \sigma_4} 2^{-4} (e^{-\beta^* \sigma_1 \sigma_4} - 1) (e^{-\beta^* \sigma_4 \sigma_3} - 1) (e^{-\beta^* \sigma_3 \sigma_2} - 1) (e^{-\beta^* \sigma_2 \sigma_1} - 1) \\ &= -\beta^{*4} + O(\beta^{*6}) \end{aligned}$$

while

$$\zeta^0(f) = +\beta^{*4} + O(\beta^{*6})$$

hence

$$Z_\Lambda / Z_\Lambda^0 = \exp(-\beta^{*4}(4L) + O(\beta^{*6})L)$$

since also the contribution for the more complicated Γ 's must be at least of order β^{*6} .

It seems clear to us that the formula that Wegner gives without proof in his work has been obtained by an expansion of the above type which is a very familiar expansion technique for the physicists.

References:

- 1) E. Wegner: J. Math. Phys. 12, 2259, 1971.
- 2) C. Gruber, H. Kunz: Comm. Math. Phys. 22, 133, 1971.

This paper has been published in *Mathematical Problems in Theoretical Physics* (Proceedings of the Conference of the International Association of Mathematical Physics, held in Rome, June 6–15, 1977), edited by G.F. Dell'Antonio, S. Doplicher and G. Jona-Lasinio, Lectures Notes in Physics no. 80, Springer, Berlin, 1978 (ISBN 3-540-08853-9), pp. 436–438.

The talk by E. Seiler, entitled *Lattice Gauge Theories*, appears in the same volume, pp. 26–36.